

BIVARIATE FREE KNOT SPLINES*

TORSTEN SCHÜTZE¹ and HUBERT SCHWETLICK²[†]

¹Siemens AG, Corporate Technology, Information & Communications, Security (CT IC 3), D-81730 Munich, Germany. torsten.schuetze@mchp.siemens.de, http://www.math.tu-dresden.de/~schuetze/ ²Department of Mathematics, Dresden University of Technology, D-01062 Dresden, Germany. schwetlick@math.tu-dresden.de, http://www.math.tu-dresden.de/~schwetli/

Abstract.

We consider the least squares approximation of gridded 2D data by tensor product splines with free knots. The smoothing functional to be minimized—a generalization of the univariate Schoenberg functional—is chosen in such a way that the solution of the bivariate problem separates into the solution of a sequence of univariate problems in case of fixed knots. The resulting optimization problem is a constrained separable least squares problem with tensor product structure. Based on some ideas developed by the authors for the univariate case, an efficient method for solving the specially structured 2D problem is proposed, analyzed and tested on hand of some examples from the literature.

AMS subject classification: Primary 65D10, 65D07; Secondary 41A15, 65D17.

Key words: Data fitting, bivariate least squares approximation, splines with free knots, constrained separable optimization problems.

1 Introduction.

Approximation by tensor product splines with fixed knots is of great importance in theory and applications, mainly due its simple structure and its ease of implementation. However, it is well known that the approximation error is, in general, much smaller if variable knots are allowed instead of fixed ones. This is true for the univariate as well as for the bivariate case.

On the other hand, approximation by splines with free knots leads to difficult but highly structured nonlinear optimization problems. There is a vast amount of literature on Chebyshev approximation by splines with free knots. Up to know mainly univariate problems have been considered. However, a few years ago Nürnberger [21] stated some research problems concerned with bivariate splines with free knots, and in [19] an algorithm was given for bivariate segment approximation which lead to good Chebyshev approximations by splines with free knots.

^{*}Received September 20, 2001. Revised January 15, 2002. Communicated by Tom Lyche. $^\dagger Research$ of the first author was partly supported by Deutsche Forschungsgemeinschaft under grant GR 705/4-2

Unlike the authors mentioned above we, however, address in this paper the problem of *least squares approximation* by splines with free knots. We generalize results from least squares approximation by univariate splines with free knots to the case of bivariate tensor product splines with gridded data. While there exist a number of papers for univariate splines with free knots, see [27] for unconstrained free knot splines, [25] and the references cited therein for constrained free knot splines, this idea seems to be new for bivariate tensor product splines.

This paper is organized as follows: In Section 2 we introduce the notation and review some results from the case of univariate spline approximation and smoothing with free knots. In the following Section 3 we consider bivariate smoothing splines and formulate the full and reduced approximation and smoothing problem, respectively. The full problems are nonlinear least squares problems with a special structure—so-called separable least squares problems with tensor product structure. In Section 4 we examine general problems of this type and show that the full and the reduced problem are equivalent in a certain sense. We apply these techniques to bivariate tensor product splines with free knots in Section 5. In Section 6 the numerical solution of the reduced problem is considered. Finally, in Section 7 we demonstrate the performance and the advantages of our method by some numerical tests.

$\mathbf{2}$ Review of univariate results.

In this section we introduce the notation and shortly review the main results of univariate free knot spline approximation. For details, discussions, and proofs we refer to [27] and [25].

2.1Spline smoothing with fixed knots.

We want to approximate given noisy data $\{x_i, y_i\}$ (i = 1, ..., m) by a function s from $\mathcal{S}_{k,\tau}$, the *n*-dimensional linear space of polynomial splines of order $k \geq 1$ with knot sequence $\tau \in \mathbb{R}^{n+k}$, where $m \ge n$. The abscissae x_i are monotonically increasing, $a \leq x_1 < \cdots < x_m \leq b$, and the noisy measurements $y_i = g(x_i) + \varepsilon_i$ (i = 1, ..., m) result from an unknown smooth function $g \in W_2^q[a, b]$ with ε_i being independent and identically distributed stochastic errors.

The parameters of the spline s have to be chosen in such a way that the Schoenberg functional

$$\frac{1}{2}\sum_{i=1}^{m} \left[y_i - s(x_i)\right]^2 + \mu \frac{1}{2} \int_a^b \left[s^{(r)}(x)\right]^2 dx$$

with the smoothing parameter $\mu > 0$ and fixed order $r \in \{0, \ldots, q\}$ in the smoothing term becomes minimal. Let $\boldsymbol{\tau} = (\tau_1, \dots, \tau_{n+k})^T$ with

 $\tau_1 = \dots = \tau_k = a < \tau_{k+1} \le \dots \le \tau_n < b = \tau_{n+1} = \dots = \tau_{n+k}$

be the knot sequence and consider a spline $s \in \mathcal{S}_{k,\tau}$ defined by $s = \sum_{j=1}^{n} B_{j,k,\tau} \alpha_j$ where $B_{j,k,\tau}$ denotes the usual *j*-th normalized polynomial B-spline of order k with knot sequence τ . Using the observation matrix

$$\mathbf{B}(\boldsymbol{\tau}) := (B_{j,k,\boldsymbol{\tau}}(x_i))_{i=1,\dots,m}^{j=1,\dots,n}, \quad \mathbf{B}(\boldsymbol{\tau}) \in \mathbb{R}^{m \times n},$$

the vector $\mathbf{y} := (y_1, \ldots, y_m)^T \in \mathbb{R}^m$ of data and the vector $\boldsymbol{\alpha} := (\alpha_1, \ldots, \alpha_n)^T \in \mathbb{R}^n$ of spline coefficients, the *approximation term* can be written as

$$\frac{1}{2}\sum_{i=1}^{m} [y_i - s(x_i)]^2 = \frac{1}{2} \|\mathbf{y} - \mathbf{B}(\boldsymbol{\tau})\boldsymbol{\alpha}\|_2^2.$$

The smoothing term can be represented in a similar form as

$$\frac{1}{2} \int_{a}^{b} \left[s^{(r)}(x) \right]^{2} dx = \frac{1}{2} \left\| \mathbf{S}_{r}(\boldsymbol{\tau}) \boldsymbol{\alpha} \right\|_{2}^{2}$$

with the smoothing matrix $\mathbf{S}_r(\boldsymbol{\tau}) \in \mathbb{R}^{(n-r) \times n}$ which is either the exact smoothing matrix $\bar{\mathbf{S}}_r$ defined as above or else a cheaper approximation $\tilde{\mathbf{S}}_r$ given in [26], [27].

Now we are able to express the Schoenberg functional as function of the spline coefficients α and the knots τ as

$$\frac{1}{2} \|\mathbf{y} - \mathbf{B}(\boldsymbol{\tau})\boldsymbol{\alpha}\|_{2}^{2} + \frac{1}{2}\mu \|\mathbf{S}_{r}(\boldsymbol{\tau})\boldsymbol{\alpha}\|_{2}^{2} = \frac{1}{2} \left\| \begin{pmatrix} \mathbf{y} \\ \mathbf{0} \end{pmatrix} - \begin{bmatrix} \mathbf{B}(\boldsymbol{\tau}) \\ \sqrt{\mu}\mathbf{S}_{r}(\boldsymbol{\tau}) \end{bmatrix} \boldsymbol{\alpha} \right\|_{2}^{2}.$$

The system matrix $\mathbf{B}_{\mu}(\boldsymbol{\tau}) := \begin{bmatrix} \mathbf{B}(\boldsymbol{\tau}) \\ \sqrt{\mu} \mathbf{S}_{r}(\boldsymbol{\tau}) \end{bmatrix} \in \mathbb{R}^{(m+n-r) \times n}$ has full rank n if the regularity condition $m \ge r$ and $\mu > 0$ is met.

2.2 Spline smoothing with free knots.

We include a subset $\mathbf{t} = (\tau_{p(1)}, \ldots, \tau_{p(l)})^T \in \mathbb{R}^l$ of the inner knots, the so-called *free knots*, into the optimization process whereas the remaining knots have to be given in advance and stay fixed. Hence, \mathbf{B} and \mathbf{S}_r become functions of \mathbf{t} alone, and we write $\mathbf{B}(\mathbf{t})$ and $\mathbf{S}_r(\mathbf{t})$ instead of $\mathbf{B}(\boldsymbol{\tau})$ and $\mathbf{S}_r(\boldsymbol{\tau})$, respectively. Note that the number l of the free knots $\tau_{p(j)}$ and the indices p(j) of them have to satisfy $l \leq n-k$ and $k < p(1) < \cdots < p(l) < n+1$.

In approximation by splines with free knots one has to avoid the coalescing of knots. Thus we require $\tau_{p(j)} \in [\tau_{p(j)-1} + \epsilon \delta_j, \tau_{p(j)+1} - \epsilon \delta_j], \, \delta_j := \tau_{p(j)+1} - \tau_{p(j)-1}$ $(j = 1, \ldots, l)$ with $0 < \epsilon \ll 1$ which, by using appropriately chosen **C** and **h**, can equivalently be written as

(2.1)
$$\mathbf{Ct} - \mathbf{h} \ge \mathbf{0} \text{ with } \mathbf{C} \in \mathbb{R}^{2l \times l}, \mathbf{h} \in \mathbb{R}^{2l}.$$

Finally we can formulate the *full smoothing problem*

(2.2) minimize
$$\mathfrak{f}(\boldsymbol{\alpha}, \mathbf{t}) := \frac{1}{2} \left\| \begin{pmatrix} \mathbf{y} \\ \mathbf{0} \end{pmatrix} - \begin{bmatrix} \mathbf{B}(\mathbf{t}) \\ \sqrt{\mu} \mathbf{S}_r(\mathbf{t}) \end{bmatrix} \boldsymbol{\alpha} \right\|_2^2 \quad \text{s.t. } \mathbf{C}\mathbf{t} - \mathbf{h} \ge \mathbf{0}.$$

Problem (2.2) is a separable nonlinear least squares problem. By inserting the minimum norm solution $\alpha_{opt}(\mathbf{t}) := \mathbf{B}_{\mu}(\mathbf{t})^+ \begin{pmatrix} \mathbf{y} \\ \mathbf{0} \end{pmatrix}$ of (2.2) for fixed knots \mathbf{t} with

respect to $\pmb{\alpha}$ into the objective function of the full problem we obtain the reduced smoothing problem

(2.3)

$$\underset{\mathbf{t}\in\mathbb{R}^{l}}{\operatorname{minimize}} f(\mathbf{t}) := \frac{1}{2} \left\| \begin{pmatrix} \mathbf{y} \\ \mathbf{0} \end{pmatrix} - \begin{bmatrix} \mathbf{B}(\mathbf{t}) \\ \sqrt{\mu} \mathbf{S}_{r}(\mathbf{t}) \end{bmatrix} \begin{bmatrix} \mathbf{B}(\mathbf{t}) \\ \sqrt{\mu} \mathbf{S}_{r}(\mathbf{t}) \end{bmatrix}^{+} \begin{pmatrix} \mathbf{y} \\ \mathbf{0} \end{pmatrix} \right\|_{2}^{2} \quad \text{s.t. } \mathbf{C}\mathbf{t} - \mathbf{h} \ge \mathbf{0}$$

with only the free knots \mathbf{t} as variables. The investigation of the full and reduced smoothing problem and the efficient numerical solution of the reduced problem was the subject of [27]. There it was shown that the reduced smoothing problem (2.3) has a solution under the conditions

- (C1) The knots satisfy $\tau_j < \tau_{j+k-q}$ $(j = q+1, \ldots, n)$.
- (C2) The regularity condition $m \ge r$ and $\mu > 0$ is met.

If, further, the condition

(C3) The free knots are simple knots, and it holds $k \geq 3$.

is fulfilled, then the change from minimizing the full functional to minimizing the reduced functional does not add any critical points, does not exclude the solution of the original problem, and the critical points are equivalent, see the original papers for details.

The reduced problem was solved by a generalized Gauss–Newton method which requires the Jacobian $\mathbf{F}'(\mathbf{t})$ of the reduced functional $f(\mathbf{t}) = \frac{1}{2} ||\mathbf{F}(\mathbf{t})||_2^2$. Since the computation of the Jacobian is quite expensive we use instead the so-called *Kaufman approximation* $\mathbf{J}_K(\mathbf{t}) \approx \mathbf{F}'(\mathbf{t})$. This results in a very efficient and robust algorithm for the computation of univariate free knot splines.

In the following we will extend these ideas to two dimensions, i.e., we tensorize the univariate results. For simplicity we use the notations from the univariate case and characterize the variables in x- and y-direction by a sub- or superscript 1 and 2, resp., e.g., τ^1 denotes the knot sequence in x-direction etc.

3 Bivariate smoothing splines.

Let $\mathbf{Z} = \{z_{i_1,i_2} : i_1 = 1, \dots, m_1; i_2 = 1, \dots, m_2\}$ be noisy measurements of an unknown function $g \in W_2^{q_1,q_2}([a_1,b_1] \times [a_2,b_2])$, which are given on a grid $[x_1,\dots,x_{m_1}] \times [y_1,\dots,y_{m_2}]$, i.e., it holds

$$z_{i_1,i_2} = g(x_{i_1}, y_{i_2}) + \varepsilon_{i_1,i_2}$$

with the abscissae $a_1 \leq x_1 < \cdots < x_{m_1} \leq b_1$, $a_2 \leq y_1 < \cdots < y_{m_2} \leq b_2$ and the measurement errors ε_{i_1,i_2} . The stochastic errors ε_{i_1,i_2} are assumed to be independent and identically distributed.

We want to approximate these data by a tensor product spline. Such splines have a simple structure, and their computation allows the separation into a sequence of univariate problems if the data are given on a rectangular grid as in our case. The extremal properties of univariate splines carry over to both interpolating and smoothing tensor product splines. For example, the natural smoothing bicubic spline is the solution of the variational problem

(3.1)
$$\min\left\{\sum_{i_1=1}^{m_1}\sum_{i_2=1}^{m_2} [z_{i_1,i_2} - s(x_{i_1}, y_{i_2})]^2 + \mu_1 \sum_{i_2=1}^{m_2} \int_{a_1}^{b_1} [D^{2,0}s(x, y_{i_2})]^2 dx + \mu_2 \sum_{i_1=1}^{m_1} \int_{a_2}^{b_2} [D^{0,2}s(x_{i_1}, y)]^2 dy + \mu_1 \mu_2 \int_{a_1}^{b_1} \int_{a_2}^{b_2} [D^{2,2}s(x, y)]^2 dx dy\right\}$$

over $s \in W_2^{2,2}([a_1, b_1] \times [a_2, b_2])$, see [15]. Here the operator D^{r_1, r_2} denotes the partial derivative of order r_1 with respect to x and of order r_2 with respect to y, and the parameters $\mu_1 > 0$ and $\mu_2 > 0$ are the smoothing parameters.

Splines defined by the above extremal property have the disadvantage that the knots of the spline are identical with the data points, i.e., no data reduction is possible. Therefore we do not use the classical variational approach but follow a so-called direct approach by restricting the approximating functions s from $W_2^{2,2}$ a priori to a fixed tensor product space of B-splines, i.e., the unknown function g will be approximated by a bivariate spline $s \in S_{k_1,\tau^1} \otimes S_{k_2,\tau^2}$ with knot sequences the cardinality of which may be much smaller than the cardinality of the grid points. The space $S_{k_1,\tau^1} \otimes S_{k_2,\tau^2}$ is the tensor product of the univariate spline spaces S_{k_1,τ^1} and S_{k_2,τ^2} of polynomial splines of order k_1 and k_2 with knot sequence τ^1 and τ^2 , respectively. The knot sequences are as follows:

$$\boldsymbol{\tau}^{1}: \tau_{1}^{1} = \ldots = \tau_{k_{1}}^{1} = a_{1} < \tau_{k_{1}+1}^{1} \leq \ldots \leq \tau_{n_{1}}^{1} < b_{1} = \tau_{n_{1}+1}^{1} = \ldots = \tau_{n_{1}+k_{1}}^{1},$$
$$\boldsymbol{\tau}^{2}: \tau_{1}^{2} = \ldots = \tau_{k_{2}}^{2} = a_{2} < \tau_{k_{2}+1}^{2} \leq \ldots \leq \tau_{n_{2}}^{2} < b_{2} = \tau_{n_{2}+1}^{2} = \ldots = \tau_{n_{2}+k_{2}}^{2}.$$

The parameters of the spline s have to be chosen in such a way that the least squares error defined by the approximation term

(3.2a)
$$\varphi(s) := \frac{1}{2} \sum_{i_1=1}^{m_1} \sum_{i_2=1}^{m_2} \left[z_{i_1,i_2} - s(x_{i_1}, y_{i_2}) \right]^2$$

becomes minimal. It is known that the solution of (3.2a) is unique if the Schoenberg-Whitney condition is fulfilled but this can not be assured in case of arbitrary data $\{x_{i_1}, y_{i_2}\}$.

By using the thin plate functional

.

$$\int_{a_1}^{b_1} \int_{a_2}^{b_2} \{ [D^{2,0}s(x,y)]^2 + 2[D^{1,1}s(x,y)]^2 + [D^{0,2}s(x,y)]^2 \} \, dx \, dy$$

as smoothing term uniqueness can be achieved independently of the data. This functional has the drawback that it has no tensor product structure and, therefore, the solution process does not separate in the case of gridded data. However, by utilizing a smoothing term which is slightly modified compared to (3.1) and minimizing the functional ϕ defined by

$$(3.2b)$$

$$\phi(s) := \frac{1}{2} \sum_{i_1=1}^{m_1} \sum_{i_2=1}^{m_2} \left[z_{i_1,i_2} - s(x_{i_1}, y_{i_2}) \right]^2 + \mu_1 \frac{1}{2} \sum_{i_2=1}^{m_2} \int_{a_1}^{b_1} \left[D^{r_1,0} s(x, y_{i_2}) \right]^2 \, dx$$

$$+ \mu_2 \frac{1}{2} \sum_{i_1=1}^{m_1} \int_{a_2}^{b_2} \left[D^{0,r_2} s(x_{i_1}, y) \right]^2 \, dy + \mu_1 \mu_2 \frac{1}{2} \int_{a_1}^{b_1} \int_{a_2}^{b_2} \left[D^{r_1,r_2} s(x, y) \right]^2 \, dy \, dx,$$

we again have a separation into a sequence of univariate problems. Unlike the thin plate functional, this smoothing functional—a generalization of the univariate Schoenberg functional—has no nice physical interpretation. However, it serves quite well as an appropriate regularization term in that it guarantees a unique solution and preserves the tensor product structure.

Bivariate smoothing splines with fixed knots have a long history: Dierckx [6] used first a nonseparable smoothing term before a suitable separable term had been found in [7]. Based on a variational approach, Hu/Schumaker considered natural bicubic smoothing splines [15] and complete smoothing splines [16]. The abstract case of interpolating and smoothing tensor product splines—from which many of the above can be derived as special cases—is investigated in [9]. Finally, in [20], [3] and [22] numerical techniques for minimizing the functional (3.2b) for fixed knots have been considered. Let us point out that, independent of the cited sources, V. Kunert has proposed such a tensor product smoothing term and, unlike the others, developed an efficient and stable solution process along the lines of [26].

The main advantage of the direct approach is that data reduction is possible and that the number and position of knots can be chosen independent of the data.

3.1 Representation of the smoothing functional

We use polynomial B-splines of order k_1 and k_2 with knot sequences τ^1 and τ^2 as basis for the univariate spaces S_{k_1,τ^1} and S_{k_2,τ^2} , respectively. They are denoted by B_{j_1,k_1,τ^1} $(j_1 = 1, \ldots, n_1)$ and B_{j_2,k_2,τ^2} $(j_2 = 1, \ldots, n_2)$. So one obtains the representation

$$s(x,y) = \sum_{j_1=1}^{n_1} \sum_{j_2=1}^{n_2} B_{j_1,k_1,\tau^1}(x) B_{j_2,k_2,\tau^2}(y) \alpha_{j_1,j_2}$$

with coefficients α_{j_1,j_2} for a tensor product spline $s \in \mathcal{S}_{k_1,\tau^1} \otimes \mathcal{S}_{k_2,\tau^2}$. By using this representation, the functionals (3.2a) and (3.2b) lead to the problems

$$(3.3a) \quad \frac{1}{2} \sum_{i_1=1}^{m_1} \sum_{i_2=1}^{m_2} \left[z_{i_1,i_2} - \sum_{j_1=1}^{n_1} \sum_{j_2=1}^{n_2} B_{j_1,k_1,\tau^1}(x_{i_1}) B_{j_2,k_2,\tau^2}(y_{i_2}) \alpha_{j_1,j_2} \right]^2 \to \min_{\alpha_{j_1,j_2}} \left[z_{i_1,i_2} - \sum_{j_1=1}^{n_1} \sum_{j_2=1}^{n_2} B_{j_1,k_1,\tau^1}(x_{i_1}) B_{j_2,k_2,\tau^2}(y_{i_2}) \alpha_{j_1,j_2} \right]^2 \to \min_{\alpha_{j_1,j_2}} \left[z_{i_1,i_2} - \sum_{j_1=1}^{n_1} \sum_{j_2=1}^{n_2} B_{j_1,k_1,\tau^1}(x_{i_1}) B_{j_2,k_2,\tau^2}(y_{i_2}) \alpha_{j_1,j_2} \right]^2 \to \min_{\alpha_{j_1,j_2}} \left[z_{i_1,i_2} - \sum_{j_1=1}^{n_1} \sum_{j_2=1}^{n_2} B_{j_1,k_1,\tau^1}(x_{i_1}) B_{j_2,k_2,\tau^2}(y_{i_2}) \alpha_{j_1,j_2} \right]^2 \to \min_{\alpha_{j_1,j_2}} \left[z_{i_1,i_2} - \sum_{j_1=1}^{n_1} \sum_{j_2=1}^{n_2} B_{j_1,k_1,\tau^1}(x_{i_1}) B_{j_2,k_2,\tau^2}(y_{i_2}) \alpha_{j_1,j_2} \right]^2 \right]^2$$

$$(3.3b) \quad \frac{1}{2} \sum_{i_{1}=1}^{m_{1}} \sum_{i_{2}=1}^{m_{2}} \left[z_{i_{1},i_{2}} - \sum_{j_{1}=1}^{n_{1}} \sum_{j_{2}=1}^{n_{2}} B_{j_{1},k_{1},\tau^{1}}(x_{i_{1}}) B_{j_{2},k_{2},\tau^{2}}(y_{i_{2}}) \alpha_{j_{1},j_{2}} \right]^{2} \\ + \mu_{1} \frac{1}{2} \sum_{i_{2}=1}^{m_{2}} \int_{a_{1}}^{b_{1}} \left[\sum_{j_{1}=1}^{n_{1}} \sum_{j_{2}=1}^{n_{2}} B_{j_{1},k_{1},\tau^{1}}(x) B_{j_{2},k_{2},\tau^{2}}(y_{i_{2}}) \alpha_{j_{1},j_{2}} \right]^{2} dx \\ + \mu_{2} \frac{1}{2} \sum_{i_{1}=1}^{m_{1}} \int_{a_{2}}^{b_{2}} \left[\sum_{j_{1}=1}^{n_{1}} \sum_{j_{2}=1}^{n_{2}} B_{j_{1},k_{1},\tau^{1}}(x_{i_{1}}) B_{j_{2},k_{2},\tau^{2}}(y) \alpha_{j_{1},j_{2}} \right]^{2} dy \\ + \mu_{1} \mu_{2} \frac{1}{2} \int_{a_{1}}^{b_{1}} \int_{a_{2}}^{b_{2}} \left[B_{j_{1},k_{1},\tau^{1}}(x) B_{j_{2},k_{2},\tau^{2}}(y) \alpha_{j_{1},j_{2}} \right]^{2} dy dx \to \min_{\alpha_{j_{1},j_{2}}}.$$

3.1.1 Matrix formulation.

For notational convenience we will use matrix notation in the following. We define

$$\mathbf{A} := (\alpha_{j_1,j_2})_{j_1=1,\dots,n_1}^{j_2=1,\dots,n_2} \in \mathbb{R}^{n_1 \times n_2}, \qquad \mathbf{Z} := (z_{i_1,i_2})_{i_1=1,\dots,m_1}^{i_2=1,\dots,m_2} \in \mathbb{R}^{m_1 \times m_2},$$

$$\boldsymbol{\beta}^1(x, \boldsymbol{\tau}^1) := (B_{1,k_1,\boldsymbol{\tau}^1}(x),\dots,B_{n_1,k_1,\boldsymbol{\tau}^1}(x))^T \in \mathbb{R}^{n_1},$$

$$\boldsymbol{\beta}^2(y, \boldsymbol{\tau}^2) := (B_{1,k_2,\boldsymbol{\tau}^2}(y),\dots,B_{n_2,k_2,\boldsymbol{\tau}^2}(y))^T \in \mathbb{R}^{n_2},$$

$$\mathbf{B}_1(\boldsymbol{\tau}^1) := (B_{j_1,k_1,\boldsymbol{\tau}^1}(x_{i_1}))_{i_1=1,\dots,m_1}^{j_1=1,\dots,m_1} = (\boldsymbol{\beta}^1(x_{i_1},\boldsymbol{\tau}^1)^T)_{i_1=1,\dots,m_1} \in \mathbb{R}^{m_1 \times n_1},$$

$$\mathbf{B}_2(\boldsymbol{\tau}^2) := (B_{j_2,k_2,\boldsymbol{\tau}^2}(y_{i_2}))_{i_2=1,\dots,m_2}^{j_2=1,\dots,m_2} = (\boldsymbol{\beta}^2(y_{i_2},\boldsymbol{\tau}^2)^T)_{i_2=1,\dots,m_2} \in \mathbb{R}^{m_2 \times n_2}.$$

Thus we have $s(x,y) = \beta^1(x, \tau^1)^T \mathbf{A} \beta^2(y, \tau^2)$, and (3.3a) reads as

(3.4a)
$$\frac{1}{2} \left\| \mathbf{Z} - \mathbf{B}_1(\boldsymbol{\tau}^1) \mathbf{A} \mathbf{B}_2(\boldsymbol{\tau}^2)^T \right\|_F^2 \to \min_{\mathbf{A} \in \mathbb{R}^{n_1 \times n_2}}$$

where $\|\,.\,\|_F$ denotes the Frobenius matrix norm. If we define smoothing matrices

$$\mathbf{S}_{r_1}^1(oldsymbol{ au}^1) \in \mathbb{R}^{(n_1-r_1) imes n_1}, \qquad \mathbf{S}_{r_2}^2(oldsymbol{ au}^2) \in \mathbb{R}^{(n_2-r_2) imes n_2}$$

as in the univariate case, (3.3b) becomes

$$\frac{1}{2} \left\| \mathbf{Z} - \mathbf{B}_{1}(\boldsymbol{\tau}^{1}) \mathbf{A} \mathbf{B}_{2}(\boldsymbol{\tau}^{2})^{T} \right\|_{F}^{2} + \frac{1}{2} \mu_{1} \left\| \mathbf{S}_{r_{1}}^{1}(\boldsymbol{\tau}^{1}) \mathbf{A} \mathbf{B}_{2}(\boldsymbol{\tau}^{2})^{T} \right\|_{F}^{2} + \frac{1}{2} \mu_{2} \left\| \mathbf{B}_{1}(\boldsymbol{\tau}^{1}) \mathbf{A} \mathbf{S}_{r_{2}}^{2}(\boldsymbol{\tau}^{2})^{T} \right\|_{F}^{2} + \frac{1}{2} \mu_{1} \mu_{2} \left\| \mathbf{S}_{r_{1}}^{1}(\boldsymbol{\tau}^{1}) \mathbf{A} \mathbf{S}_{r_{2}}^{2}(\boldsymbol{\tau}^{2})^{T} \right\|_{F}^{2} \to \min_{\mathbf{A} \in \mathbb{R}^{n_{1} \times n_{2}}}$$

or, using block notation,

(3.4b)
$$\frac{1}{2} \left\| \begin{bmatrix} \mathbf{Z} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} - \begin{bmatrix} \mathbf{B}_1(\boldsymbol{\tau}^1) \\ \sqrt{\mu_1} \mathbf{S}_{r_1}^1(\boldsymbol{\tau}^1) \end{bmatrix} \mathbf{A} \begin{bmatrix} \mathbf{B}_2(\boldsymbol{\tau}^2) \\ \sqrt{\mu_2} \mathbf{S}_{r_2}^2(\boldsymbol{\tau}^2) \end{bmatrix}^T \right\|_F^2 \to \min_{\mathbf{A} \in \mathbb{R}^{n_1 \times n_2}}.$$

Before we characterize the solutions of (3.4a) and (3.4b) we need some auxiliary results. Recall that the *Kronecker product* of $\mathbf{A} \in \mathbb{R}^{m \times n}$ and $\mathbf{B} \in \mathbb{R}^{p \times q}$ is given by

$$\mathbf{A} \otimes \mathbf{B} := \begin{bmatrix} a_{11}\mathbf{B} & \cdots & a_{1n}\mathbf{B} \\ \vdots & & \vdots \\ a_{m1}\mathbf{B} & \cdots & a_{mn}\mathbf{B} \end{bmatrix} \in \mathbb{R}^{(m \cdot p) \times (n \cdot q)}.$$

For matrices $\mathbf{A} = (a_{i,j}) \in \mathbb{R}^{m \times n}$ the operation vec : $\mathbb{R}^{m \times n} \to \mathbb{R}^{m \cdot n}$ is defined by

$$\operatorname{vec}(\mathbf{A}) := (a_{11}, \dots, a_{m1}, a_{12}, \dots, a_{m2}, \dots, a_{1n}, \dots, a_{mn})^T \in \mathbb{R}^{m \cdot n}$$

Using elementary facts about Kronecker products and the vec-operator, see e.g. [1], it can be shown that $\|\mathbf{A}\mathbf{X}\mathbf{B} - \mathbf{C}\|_{F}^{2} = \|(\mathbf{B}^{T} \otimes \mathbf{A}) \operatorname{vec}(\mathbf{X}) - \operatorname{vec}(\mathbf{C})\|_{2}^{2}$. Hence, problem (3.4a) is equivalent to

(3.5)
$$\frac{1}{2} \|\operatorname{vec}\left(\mathbf{Z}\right) - \left(\mathbf{B}_{2}(\boldsymbol{\tau}^{2}) \otimes \mathbf{B}_{1}(\boldsymbol{\tau}^{1})\right) \operatorname{vec}\left(\mathbf{A}\right)\|_{2}^{2} \to \min_{\operatorname{vec}\left(\mathbf{A}\right) \in \mathbb{R}^{n_{1}n_{2}}},$$

the minimum norm solution $\mathbf{A}_{opt}(\boldsymbol{\tau}^1, \boldsymbol{\tau}^2)$ of which is, for fixed $\boldsymbol{\tau}^1$ and $\boldsymbol{\tau}^2$, given by

$$\operatorname{vec}\left(\mathbf{A}_{opt}\right) = (\mathbf{B}_{2} \otimes \mathbf{B}_{1})^{+} \operatorname{vec}\left(\mathbf{Z}\right) = (\mathbf{B}_{2}^{+} \otimes \mathbf{B}_{1}^{+}) \operatorname{vec}\left(\mathbf{Z}\right) = \operatorname{vec}\left(\mathbf{B}_{1}^{+} \mathbf{Z} (\mathbf{B}_{2}^{+})^{T}\right), \text{i.e.},$$

(3.6a)
$$\mathbf{A}_{opt}(\boldsymbol{\tau}^1, \boldsymbol{\tau}^2) := \mathbf{B}_1(\boldsymbol{\tau}^1)^+ \mathbf{Z} \left(\mathbf{B}_2(\boldsymbol{\tau}^2)^+ \right)^T.$$

It can be computed by successive solution of the following univariate problems:

(F) Solve
$$\frac{1}{2} \|\mathbf{Z} - \mathbf{B}_1(\boldsymbol{\tau}^1)\mathbf{F}\|_F^2 \to \min_{\mathbf{F} \in \mathbb{R}^{n_1 \times m_2}} \quad \text{for} \quad \mathbf{F} = \mathbf{B}_1(\boldsymbol{\tau}^1)^+ \mathbf{Z},$$

(A) Solve
$$\frac{1}{2} \left\| \mathbf{F}^T - \mathbf{B}_2(\boldsymbol{\tau}^2) \mathbf{A}^T \right\|_F^2 \to \min_{\mathbf{A} \in \mathbb{R}^{n_1 \times n_2}} \quad \text{for} \quad \mathbf{A}^T = \mathbf{B}_2(\boldsymbol{\tau}^2)^+ \mathbf{F}^T.$$

Analogously one obtains the minimum norm solution to problem (3.4b)

(3.6b)
$$\mathbf{A}_{opt}(\boldsymbol{\tau}^1, \boldsymbol{\tau}^2) := \begin{bmatrix} \mathbf{B}_1(\boldsymbol{\tau}^1) \\ \sqrt{\mu_1} \mathbf{S}_{r_1}^1(\boldsymbol{\tau}^1) \end{bmatrix}^+ \begin{bmatrix} \mathbf{Z} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \left(\begin{bmatrix} \mathbf{B}_2(\boldsymbol{\tau}^2) \\ \sqrt{\mu_2} \mathbf{S}_{r_2}^2(\boldsymbol{\tau}^2) \end{bmatrix}^+ \right)^T$$

which can successively be computed as follows:

4

(F) Solve
$$\frac{1}{2} \left\| \begin{bmatrix} \mathbf{Z} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} - \begin{bmatrix} \mathbf{B}_1(\boldsymbol{\tau}^1) \\ \sqrt{\mu_1} \mathbf{S}_{r_1}^1(\boldsymbol{\tau}^1) \end{bmatrix} \mathbf{F} \right\|_F^2 \to \min_{\mathbf{F} \in \mathbb{R}^{n_1 \times (m_2 + n_2 - r_2)}} \quad \text{for} \quad \mathbf{F},$$

(A) Solve
$$\frac{1}{2} \left\| \mathbf{F}^T - \begin{bmatrix} \mathbf{B}_2(\boldsymbol{\tau}^2) \\ \sqrt{\mu_2} \mathbf{S}_{r_2}^2(\boldsymbol{\tau}^2) \end{bmatrix} \mathbf{A}^T \right\|_F^2 \to \min_{\mathbf{A} \in \mathbb{R}^{n_1 \times n_2}} \quad \text{for } \mathbf{A}^T.$$

For the solution of subproblems (A) and (F) we can use the row-wise Givens factorization developed for the univariate case, see [26] for details. General investigations about the exploitation of structure in large linear least squares problems based on Kronecker products can be found in [10].

3.2 The full and the reduced approximation problem.

Following our general philosophy, see [27], [25], we now include the knots of the splines into the optimization process. If we consider a subset $(\mathbf{t}^1, \mathbf{t}^2)$ of the inner knots in problem (3.4a) as variable, we get the *full approximation problem*

(3.7)
$$f(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) := \frac{1}{2} \left\| \mathbf{Z} - \mathbf{B}_1(\mathbf{t}^1) \mathbf{A} \mathbf{B}_2(\mathbf{t}^2)^T \right\|_F^2 \to \min_{\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}}$$

with linear constraints

$$(3.8) \mathbf{C}_1 \mathbf{t}^1 - \mathbf{h}^1 \ge \mathbf{0}, \quad \mathbf{C}_2 \mathbf{t}^2 - \mathbf{h}^2 \ge \mathbf{0}.$$

which avoid coalescing of knots.

Inserting the minimum norm solution (3.6a) into the functional \mathfrak{f} one obtains the reduced approximation problem

(3.9)
$$f(\mathbf{t}^1, \mathbf{t}^2) := \frac{1}{2} \left\| \mathbf{Z} - \mathbf{B}_1(\mathbf{t}^1) \mathbf{B}_1(\mathbf{t}^1)^+ \mathbf{Z} \left(\mathbf{B}_2(\mathbf{t}^2)^+ \right)^T \mathbf{B}_2(\mathbf{t}^2)^T \right\|_F^2 \to \min_{\mathbf{t}^1, \mathbf{t}^2}$$

with the linear inequality constraints (3.8). The objective function can equivalently be written as

$$f(\mathbf{t}^{1}, \mathbf{t}^{2}) = \frac{1}{2} \left\| \mathbf{Z} - \mathbf{P}_{B_{1}} \mathbf{Z} \mathbf{P}_{B_{2}} \right\|_{F}^{2} = \frac{1}{2} \left\| \mathbf{P}_{B_{1}}^{\perp} \mathbf{Z} \mathbf{P}_{B_{2}} \right\|_{F}^{2}$$

with the orthogonal projectors $\mathbf{P}_{B_1} := \mathbf{B}_1(\mathbf{t}^1)\mathbf{B}_1(\mathbf{t}^1)^+$, $\mathbf{P}_{B_2} := \mathbf{B}_2(\mathbf{t}^2)\mathbf{B}_2(\mathbf{t}^2)^+$, and $\mathbf{P}_{B_1}^{\perp} := \mathbf{I} - \mathbf{P}_{B_1}$.

3.3 The full and the reduced smoothing problem.

For the smoothing problem (3.4b) one obtains analogously the full smoothing problem

(3.10)

$$\mathfrak{f}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) := \frac{1}{2} \left\| \begin{bmatrix} \mathbf{Z} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} - \begin{bmatrix} \mathbf{B}_1(\mathbf{t}^1) \\ \sqrt{\mu_1} \mathbf{S}_{r_1}^1(\mathbf{t}^1) \end{bmatrix} \mathbf{A} \begin{bmatrix} \mathbf{B}_2(\mathbf{t}^2) \\ \sqrt{\mu_2} \mathbf{S}_{r_2}^2(\mathbf{t}^2) \end{bmatrix}^T \right\|_F^2 \to \min_{\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}}$$

with the linear inequality constraints (3.8).

Again, inserting the minimum norm solution (3.6b) into the functional, one obtains the *reduced smoothing problem*

(3.11)
$$f(\mathbf{t}^1, \mathbf{t}^2) := \frac{1}{2} \left\| \begin{bmatrix} \mathbf{Z} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} - \mathbf{P}_{\begin{bmatrix} B \\ \sqrt{\mu}S \end{bmatrix} 1} \begin{bmatrix} \mathbf{Z} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{P}_{\begin{bmatrix} B \\ \sqrt{\mu}S \end{bmatrix} 2} \right\|_F^2 \to \min_{\mathbf{t}^1, \mathbf{t}^2}$$

with the constraints (3.8). The orthogonal projectors are defined by

$$\begin{split} \mathbf{P}_{\begin{bmatrix} B\\ \sqrt{\mu}S \end{bmatrix}^1} &:= \begin{bmatrix} \mathbf{B}_1(\mathbf{t}^1)\\ \sqrt{\mu_1} \mathbf{S}_{r_1}^1(\mathbf{t}^1) \end{bmatrix} \begin{bmatrix} \mathbf{B}_1(\mathbf{t}^1)\\ \sqrt{\mu_1} \mathbf{S}_{r_1}^1(\mathbf{t}^1) \end{bmatrix}^+ \\ \mathbf{P}_{\begin{bmatrix} B\\ \sqrt{\mu}S \end{bmatrix}^2} &:= \begin{bmatrix} \mathbf{B}_2(\mathbf{t}^2)\\ \sqrt{\mu_2} \mathbf{S}_{r_2}^2(\mathbf{t}^2) \end{bmatrix} \begin{bmatrix} \mathbf{B}_2(\mathbf{t}^2)\\ \sqrt{\mu_2} \mathbf{S}_{r_2}^2(\mathbf{t}^2) \end{bmatrix}^+. \end{split}$$

In the rest of this paper we investigate the equivalence of solutions to the full and the reduced problem and develop an algorithm for the solution of the reduced problem. We are especially interested whether the problems are separable and how techniques from the univariate case carry over. Note that a satisfying resolution of the geometric structure expressed by the data is, in principle, impossible if we use tensor product splines. For example, peaks and diagonally placed layers can not be approximated well. Alternatives are splines on triangulations, curved knot lines, or the usage of hierarchical B-Splines. Despite these inherent disadvantages, tensor product splines are widely used in practice since they are simple to handle and inexpensive to compute as long as the knot lines are fixed.

While there are several algorithms available for the direct minimization of the least squares error as function of the free knots in the univariate case, we know of no such algorithms in the bivariate case. If we restrict to a heuristic, adaptive insertion of knot lines we have the algorithm REGRID from [8] at our disposal. For the case of Chebyshev approximation first results were achieved recently in [19] as already mentioned in the introduction.

4 Separable least squares problems with tensor product structure.

In this section we will detach from the problem of spline smoothing and consider general optimization problems of the following form:

Full problem

(4.1)
$$f(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) := \frac{1}{2} \left\| \mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) \right\|_F^2 \to \min_{\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}}$$

with $\mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) := \mathbf{Z} - \mathbf{B}_1(\mathbf{t}^1) \mathbf{A} \mathbf{B}_2(\mathbf{t}^2)^T$ subject to the constraints

(4.2)
$$\mathbf{C}_1 \mathbf{t}^1 - \mathbf{h}^1 \ge \mathbf{0}, \quad \mathbf{C}_2 \mathbf{t}^2 - \mathbf{h}^2 \ge \mathbf{0}.$$

Here \mathbf{B}_1 and \mathbf{B}_2 are arbitrary smooth matrix functions. The remaining quantities \mathbf{Z} , \mathbf{C}_1 , \mathbf{C}_2 , \mathbf{h}^1 and \mathbf{h}^2 are constant matrices and vectors. The variable projection solution, i. e., the optimal solution \mathbf{A}_{opt} of (4.1) for fixed \mathbf{t}^1 and \mathbf{t}^2 , is defined by

(4.3)
$$\mathbf{A}_{opt}(\mathbf{t}^1, \mathbf{t}^2) := \mathbf{B}_1(\mathbf{t}^1)^+ \mathbf{Z} \left(\mathbf{B}_2(\mathbf{t}^2)^+ \right)^T.$$

Reduced problem

(4.4)
$$f(\mathbf{t}^1, \mathbf{t}^2) := \frac{1}{2} \left\| \mathbf{F}(\mathbf{t}^1, \mathbf{t}^2) \right\|_F^2 \to \min_{\mathbf{t}^1, \mathbf{t}^2}$$

with $\mathbf{F}(\mathbf{t}^1, \mathbf{t}^2) := \mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}_{opt}(\mathbf{t}^1, \mathbf{t}^2)) = \mathbf{P}_{B_1}^{\perp} \mathbf{Z} \mathbf{P}_{B_2}$ under the constraints (4.2). Obviously an equivalent representation of the reduced functional $f(\mathbf{t}^1, \mathbf{t}^2)$ is given by

$$f(\mathbf{t}^{1},\mathbf{t}^{2}) = \frac{1}{2} \left\| \mathbf{B}_{1}(\mathbf{t}^{1})\mathbf{B}_{1}(\mathbf{t}^{1})^{+}\mathbf{Z} \left(\mathbf{B}_{2}(\mathbf{t}^{2})^{+} \right)^{T} \mathbf{B}_{2}(\mathbf{t}^{2})^{T} - \mathbf{Z} \right\|_{F}^{2} = \frac{1}{2} \left\| \mathbf{P}_{B_{1}}\mathbf{Z}\mathbf{P}_{B_{2}}^{\perp} \right\|_{F}^{2}$$

We will now—in generalization of the results of [14]—investigate to what extent the full and reduced problem are equivalent. In the following we will often need the Fréchet derivatives of certain functionals which are defined using Frobenius norms. Let ∂ denote the operator of Fréchet derivative. Then we have

LEMMA 4.1 (FRÉCHET DERIVATIVES OF FROBENIUS NORMS).

Let $\mathbf{A} : \mathbb{R}^l \to \mathfrak{L}(\mathbb{R}^n, \mathbb{R}^m)$, $\mathbf{x} \in \mathbb{R}^l \to \mathbf{A}(\mathbf{x}) \in \mathbb{R}^{m \times n}$ be a Fréchet differentiable matrix function, and define $f : \mathbb{R}^l \to \mathfrak{L}(\mathbb{R})$ by

$$\mathbf{x} \in \mathbb{R}^l \to f(\mathbf{x}) := \frac{1}{2} \|\mathbf{A}(\mathbf{x})\|_F^2 = \frac{1}{2} \operatorname{tr} \left\{ \mathbf{A}(\mathbf{x})^T \mathbf{A}(\mathbf{x}) \right\} = \frac{1}{2} \operatorname{tr} \left\{ \mathbf{A}(\mathbf{x}) \mathbf{A}(\mathbf{x})^T \right\} \in \mathbb{R}.$$

Then it holds

$$\partial f(\mathbf{x})[\Delta \mathbf{x}] = \operatorname{tr}\left\{ (\partial \mathbf{A}(\mathbf{x})[\Delta \mathbf{x}])^T \mathbf{A}(\mathbf{x}) \right\} = \operatorname{tr}\left\{ \mathbf{A}(\mathbf{x})^T (\partial \mathbf{A}(\mathbf{x})[\Delta \mathbf{x}]) \right\}$$
$$= \operatorname{tr}\left\{ (\partial \mathbf{A}(\mathbf{x})[\Delta \mathbf{x}]) \mathbf{A}(\mathbf{x})^T \right\} = \operatorname{tr}\left\{ \mathbf{A}(\mathbf{x}) (\partial \mathbf{A}(\mathbf{x})[\Delta \mathbf{x}])^T \right\}$$

for all $\Delta \mathbf{x} \in \mathbb{R}^l$.

Further, we need the Fréchet derivative of an orthogonal projector. We consider an arbitrary differentiable $m \times n$ matrix function $\mathbf{B}(.)$ of locally constant rank and the corresponding orthogonal projector $\mathbf{P}_B := \mathbf{B}\mathbf{B}^-$.

LEMMA 4.2 (GOLUB/PEREYRA [14, LEMMA 4.1]).

Let $\mathbf{B}^{-}(.)$ be an $n \times m$ matrix function such that (P1) $\mathbf{B}\mathbf{B}^{-}\mathbf{B} = \mathbf{B}$ and (P3) $(\mathbf{B}\mathbf{B}^{-})^{T} = \mathbf{B}\mathbf{B}^{-}$. Then it holds

$$\partial \mathbf{P}_B = \mathbf{P}_B^{\perp}(\partial \mathbf{B})\mathbf{B}^- + \left(\mathbf{P}_B^{\perp}(\partial \mathbf{B})\mathbf{B}^-\right)^T \quad where \quad \mathbf{P}_B^{\perp} = \mathbf{I} - \mathbf{P}_B.$$

4.1 Fréchet derivative of the full functional.

Let ∂_1 be the Fréchet derivative with respect to t^1 . Using Lemma 4.1 we have

$$\boldsymbol{\partial}_{1}\mathfrak{f}(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A})[\Delta\mathbf{t}^{1}] = \mathrm{tr}\left\{\left(\boldsymbol{\partial}_{1}\mathfrak{F}(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A})[\Delta\mathbf{t}^{1}]\right)^{T}\mathfrak{F}(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A})\right\}$$

For simplicity we will omit the argument of the matrix functions \mathbf{B}_1 and \mathbf{B}_2 . Obviously it holds $\partial_1 \mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A})[\Delta \mathbf{t}^1] = -\partial_1 \mathbf{B}_1[\Delta \mathbf{t}^1] \mathbf{A} \mathbf{B}_2^T$, hence we have

(4.5)
$$\partial_1 \mathfrak{f}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A})[\Delta \mathbf{t}^1] = -\operatorname{tr}\left\{\mathbf{B}_2 \mathbf{A}^T \left(\partial_1 \mathbf{B}_1[\Delta \mathbf{t}^1]\right)^T \left(\mathbf{Z} - \mathbf{B}_1 \mathbf{A} \mathbf{B}_2^T\right)\right\}.$$

for the Fréchet derivative of the full functional with respect to t^1 .

Let ∂_2 be the Fréchet derivative with respect to \mathbf{t}^2 . Using the representation

$$\mathfrak{f}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) = \frac{1}{2} \left\| \mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) \right\|_F^2 = \frac{1}{2} \operatorname{tr} \left\{ \mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) \mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A})^T \right\}$$

with $\mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A}) := \mathbf{B}_1 \mathbf{A} \mathbf{B}_2^T - \mathbf{Z}$ we get from Lemma 4.1

$$\partial_{2}\mathfrak{f}(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A})[\Delta\mathbf{t}^{2}] = \operatorname{tr}\left\{\mathfrak{F}(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A})\left(\partial_{2}\mathfrak{F}(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A})[\Delta\mathbf{t}^{2}]\right)^{T}\right\}.$$

With $\partial_2 \mathfrak{F}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A})[\Delta \mathbf{t}^2] = \mathbf{B}_1 \mathbf{A} \left(\partial_2 \mathbf{B}_2[\Delta \mathbf{t}^2] \right)^T$ we finally obtain

(4.6)
$$\partial_2 \mathfrak{f}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A})[\Delta \mathbf{t}^2] = \operatorname{tr}\left\{ \left(\mathbf{B}_1 \mathbf{A} \mathbf{B}_2^T - \mathbf{Z} \right) \left(\partial_2 \mathbf{B}_2[\Delta \mathbf{t}^2] \right) \mathbf{A}^T \mathbf{B}_1^T \right\}$$

for the Fréchet derivative of the full functional with respect to \mathbf{t}^2 .

4.2 Fréchet derivative of the reduced functional.

We will now compute the Fréchet derivative of the reduced functional f with respect to ${\bf t}^1.$ First, we have

$$\partial_1 f(\mathbf{t}^1, \mathbf{t}^2)[\Delta \mathbf{t}^1] = \operatorname{tr}\left\{ \left(\partial_1 \mathbf{F}(\mathbf{t}^1, \mathbf{t}^2)[\Delta \mathbf{t}^1] \right)^T \mathbf{F}(\mathbf{t}^1, \mathbf{t}^2) \right\}$$

because of Lemma 4.1, and with Lemma 4.2 $\,$

$$\begin{split} \boldsymbol{\partial}_{1}\mathbf{F}(\mathbf{t}^{1},\mathbf{t}^{2})[\Delta\mathbf{t}^{1}] &= \left(\boldsymbol{\partial}_{1}\mathbf{P}_{B_{1}}^{\perp}[\Delta\mathbf{t}^{1}]\right)\mathbf{Z}\mathbf{P}_{B_{2}} \\ &= -\left\{\mathbf{P}_{B_{1}}^{\perp}(\boldsymbol{\partial}_{1}\mathbf{B}_{1}[\Delta\mathbf{t}^{1}])\mathbf{B}_{1}^{+} + \left(\mathbf{P}_{B_{1}}^{\perp}(\boldsymbol{\partial}_{1}\mathbf{B}_{1}[\Delta\mathbf{t}^{1}])\mathbf{B}_{1}^{+}\right)^{T}\right\}\mathbf{Z}\mathbf{P}_{B_{2}}. \end{split}$$

It follows

$$\boldsymbol{\partial}_{1} f(\mathbf{t}^{1}, \mathbf{t}^{2}) [\Delta \mathbf{t}^{1}] = - \operatorname{tr} \left\{ \mathbf{P}_{B_{2}}^{T} \mathbf{Z}^{T} \left\{ \mathbf{P}_{B_{1}}^{\perp} (\boldsymbol{\partial}_{1} \mathbf{B}_{1} [\Delta \mathbf{t}^{1}]) \mathbf{B}_{1}^{+} + \left(\mathbf{P}_{B_{1}}^{\perp} (\boldsymbol{\partial}_{1} \mathbf{B}_{1} [\Delta \mathbf{t}^{1}]) \mathbf{B}_{1}^{+} \right)^{T} \right\} \mathbf{P}_{B_{1}}^{\perp} \mathbf{Z} \mathbf{P}_{B_{2}} \right\}$$

and, because of $\mathbf{B}_1^+ \mathbf{P}_{B_1}^\perp = \mathbf{0}$ and $\left(\mathbf{P}_{B_1}^\perp\right)^T \mathbf{P}_{B_1}^\perp = \mathbf{P}_{B_1}^\perp$, finally

(4.7)
$$\partial_1 f(\mathbf{t}^1, \mathbf{t}^2)[\Delta \mathbf{t}^1] = -\operatorname{tr}\left\{\mathbf{P}_{B_2}\mathbf{Z}^T\left((\partial_1 \mathbf{B}_1)[\Delta \mathbf{t}^1]\right)\mathbf{B}_1^+\right)^T \mathbf{P}_{B_1}^{\perp}\mathbf{Z}\mathbf{P}_{B_2}\right\}.$$

Analogously we obtain

(4.8)
$$\partial_2 f(\mathbf{t}^1, \mathbf{t}^2)[\Delta \mathbf{t}^2] = -\operatorname{tr}\left\{\mathbf{P}_{B_1}\mathbf{Z}\mathbf{P}_{B_2}^{\perp}(\partial_2 \mathbf{B}_2[\Delta \mathbf{t}^2])\mathbf{B}_2^{\perp}\mathbf{Z}^T\mathbf{P}_{B_1}\right\}.$$

using the representation

$$f(\mathbf{t}^{1}, \mathbf{t}^{2}) = \frac{1}{2} \|\mathbf{F}(\mathbf{t}^{1}, \mathbf{t}^{2})\|_{F}^{2} = \frac{1}{2} \operatorname{tr} \left\{ \mathbf{F}(\mathbf{t}^{1}, \mathbf{t}^{2}) \mathbf{F}(\mathbf{t}^{1}, \mathbf{t}^{2})^{T} \right\}$$

with $\mathbf{F}(\mathbf{t}^1, \mathbf{t}^2) := \mathbf{P}_{B_1} \mathbf{Z} \mathbf{P}_{B_2}^{\perp}$.

4.3 Relations between Fréchet derivatives.

LEMMA 4.3. Let the matrix functions \mathbf{B}_1 and \mathbf{B}_2 have locally constant rank around \mathbf{t}^1 and \mathbf{t}^2 , respectively, and let

$$\mathbf{A}_{opt}\left(\mathbf{t}^{1},\mathbf{t}^{2}
ight)=\mathbf{B}_{1}\left(\mathbf{t}^{1}
ight)^{+}\mathbf{Z}\left(\mathbf{B}_{2}\left(\mathbf{t}^{2}
ight)^{+}
ight)^{T}$$

be the corresponding variable projection solution. Then

$$\partial_{1}f(\mathbf{t}^{1},\mathbf{t}^{2}) = \partial_{1}\mathfrak{f}(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A}_{opt}(\mathbf{t}^{1},\mathbf{t}^{2})), \\ \partial_{2}f(\mathbf{t}^{1},\mathbf{t}^{2}) = \partial_{2}\mathfrak{f}(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A}_{opt}(\mathbf{t}^{1},\mathbf{t}^{2})).$$

PROOF. (i) First we consider the Fréchet derivative with respect to t^1 . By inserting the variable projection solution into (4.5) we obtain

$$\begin{aligned} \partial_{1}\mathfrak{f}\left(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A}_{opt}\left(\mathbf{t}^{1},\mathbf{t}^{2}\right)\right)\left[\Delta\mathbf{t}^{1}\right] \\ &= \operatorname{tr}\left\{-\mathbf{B}_{2}\mathbf{B}_{2}^{+}\mathbf{Z}^{T}(\mathbf{B}_{1}^{+})^{T}(\partial_{1}\mathbf{B}_{1}[\Delta\mathbf{t}^{1}])^{T}\left(\mathbf{Z}-\mathbf{B}_{1}\mathbf{B}_{1}^{+}\mathbf{Z}(\mathbf{B}_{2}^{+})^{T}\mathbf{B}_{2}^{T}\right)\right\} \\ &= \operatorname{tr}\left\{-\mathbf{P}_{B_{2}}\mathbf{Z}^{T}\left((\partial_{1}\mathbf{B}_{1}[\Delta\mathbf{t}^{1}])\mathbf{B}_{1}^{+}\right)^{T}\mathbf{P}_{B_{1}}^{\perp}\mathbf{Z}\mathbf{P}_{B_{2}}\right\} = \partial_{1}f\left(\mathbf{t}^{1},\mathbf{t}^{2}\right)\left[\Delta\mathbf{t}^{1}\right]. \end{aligned}$$

(ii) Analogously we obtain the Fréchet derivative with respect to \mathbf{t}^2 as

$$\begin{aligned} \boldsymbol{\partial}_{2}\mathfrak{f}\left(\mathbf{t}^{1},\mathbf{t}^{2},\mathbf{A}_{opt}\left(\mathbf{t}^{1},\mathbf{t}^{2}\right)\right)\left[\Delta\mathbf{t}^{2}\right] \\ &= \operatorname{tr}\left\{\left(\mathbf{B}_{1}\mathbf{B}_{1}^{+}\mathbf{Z}(\mathbf{B}_{2}^{+})^{T}\mathbf{B}_{2}^{T}-\mathbf{Z}\right)\left(\boldsymbol{\partial}_{2}\mathbf{B}_{2}[\Delta\mathbf{t}^{2}]\right)\mathbf{B}_{2}^{+}\mathbf{Z}^{T}(\mathbf{B}_{1}^{+})^{T}\mathbf{B}_{1}^{T}\right\} \\ &= \operatorname{tr}\left\{-\mathbf{P}_{B_{1}}\mathbf{Z}\mathbf{P}_{B_{2}}^{\perp}\left(\boldsymbol{\partial}_{2}\mathbf{B}_{2}[\Delta\mathbf{t}^{2}]\right)\mathbf{B}_{2}^{+}\mathbf{Z}^{T}\mathbf{P}_{B_{1}}\right\} = \boldsymbol{\partial}_{2}f\left(\mathbf{t}^{1},\mathbf{t}^{2}\right)[\Delta\mathbf{t}^{2}].\end{aligned}$$

The importance of the above lemma and the following theorem lies not only in their statements itself—which are to be expected—but in the formulae for gradient and Jacobian of the reduced functional derived there.

4.4 Correspondence between the full and the reduced problem.

The following theorem is a natural generalization of [14, Theorem 2.1] to the tensor product case. Note, however, that we have allowed linear inequality constraints with respect to t^1 and t^2 .

THEOREM 4.4 (CORRESPONDENCE BETWEEN THE FULL AND REDUCED PROB-LEM).

Let the full and the reduced problem be defined as above. Assume further that the matrix functions $\mathbf{B}_1(\mathbf{t}^1)$ and $\mathbf{B}_2(\mathbf{t}^2)$ have constant rank on the open sets $\Omega_1 \subset \mathbb{R}^{l_1}$ and $\Omega_2 \subset \mathbb{R}^{l_2}$, respectively.

(i) If $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$ is a critical point (or a global minimizer on $\Omega_1 \times \Omega_2$) of the reduced problem then $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}))$ is a critical point (or a global minimizer for $(\mathbf{t}^1, \mathbf{t}^2) \in \Omega_1 \times \Omega_2$) of the full problem and it holds

$$f(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*})) = f(\mathbf{t}^{1*}, \mathbf{t}^{2*})$$

where $\mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}) = \mathbf{B}_1(\mathbf{t}^{1*})^+ \mathbf{Z} (\mathbf{B}_2(\mathbf{t}^{2*})^+)^T$.

(ii) Let $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$ be a global minimizer of the full problem for $(\mathbf{t}^1, \mathbf{t}^2) \in \Omega_1 \times \Omega_2$. Then $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$ is a global minimizer of the reduced problem on $\Omega_1 \times \Omega_2$ and it holds

$$f\left(\mathbf{t}^{1*}, \mathbf{t}^{2*}\right) = \mathfrak{f}\left(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^{*}\right).$$

If there is an unique A^* among all minimizing triplets $\mathfrak{f}(t^1,t^2,A),$ then it must hold

$$\mathbf{A}^* = \mathbf{B}_1 \left(\mathbf{t}^{1*} \right)^+ \mathbf{Z} \left(\mathbf{B}_2 \left(\mathbf{t}^{2*} \right)^+ \right)^T.$$

PROOF. We define the Lagrangian of the the full and reduced problems

$$L_{I}(\mathbf{t}^{1}, \mathbf{t}^{2}, \mathbf{A}, \mathbf{w}^{1}, \mathbf{w}^{2}) := \mathfrak{f}(\mathbf{t}^{1}, \mathbf{t}^{2}, \mathbf{A}) - \sum_{i=1}^{ncstr_{1}} \mathfrak{w}_{i}^{1} r_{i}^{1}(\mathbf{t}^{1}) - \sum_{i=1}^{ncstr_{2}} \mathfrak{w}_{i}^{2} r_{i}^{2}(\mathbf{t}^{2})$$
$$L_{II}(\mathbf{t}^{1}, \mathbf{t}^{2}, \mathbf{w}^{1}, \mathbf{w}^{2}) := f(\mathbf{t}^{1}, \mathbf{t}^{2}) - \sum_{i=1}^{ncstr_{1}} w_{i}^{1} r_{i}^{1}(\mathbf{t}^{1}) - \sum_{i=1}^{ncstr_{2}} w_{i}^{2} r_{i}^{2}(\mathbf{t}^{2})$$

with nonnegative Lagrange multipliers \mathbf{w}_i^1 , w_i^1 $(i = 1, ..., ncstr_1)$ and \mathbf{w}_i^2 , w_i^2 $(i = 1, ..., ncstr_2)$; and the constraints $\mathbf{r}^1(\mathbf{t}^1) := \mathbf{C}_1\mathbf{t}^1 - \mathbf{h}^1 \ge \mathbf{0}$ and $\mathbf{r}^2(\mathbf{t}^2) := \mathbf{C}_2\mathbf{t}^2 - \mathbf{h}^2 \ge \mathbf{0}$.

(i) Let (t^{1*}, t^{2*}) be a critical point of the reduced problem. The first order optimality conditions state the existence of multipliers w^{1*} and w^{2*} so that

$$\begin{aligned} \nabla_{\mathbf{t}^{1}} L_{II}\left(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{w}^{1*}, \mathbf{w}^{2*}\right) &= \mathbf{0} & \nabla_{\mathbf{t}^{2}} L_{II}\left(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{w}^{1*}, \mathbf{w}^{2*}\right) &= \mathbf{0} \\ r_{i}^{1}(\mathbf{t}^{1*}) &\geq 0 \quad (i = 1, \dots, ncstr_{1}) & r_{i}^{2}(\mathbf{t}^{2*}) &\geq 0 \quad (i = 1, \dots, ncstr_{2}) \\ w_{i}^{1*} r_{i}^{1}(\mathbf{t}^{1*}) &= 0 \quad (i = 1, \dots, ncstr_{1}) & w_{i}^{2*} r_{i}^{2}(\mathbf{t}^{2*}) &= 0 \quad (i = 1, \dots, ncstr_{2}) \\ w_{i}^{1*} &\geq 0 \quad (i = 1, \dots, ncstr_{1}) & w_{i}^{2*} &\geq 0 \quad (i = 1, \dots, ncstr_{2}). \end{aligned}$$

Further we assume that appropriate constraint qualifications at $({\bf t}^{1*}, {\bf t}^{2*})$ are satisfied.

Now it holds

$$\mathbf{0} = \nabla_{\mathbf{t}^{1}} L_{II} \left(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{w}^{1*}, \mathbf{w}^{2*} \right)$$

= $\nabla_{\mathbf{t}^{1}} f \left(\mathbf{t}^{1*}, \mathbf{t}^{2*} \right) - \sum_{i=1}^{ncstr_{1}} w_{i}^{1*} \nabla_{\mathbf{t}^{1}} r_{i}^{1} \left(\mathbf{t}^{1*} \right) - \sum_{i=1}^{ncstr_{2}} w_{i}^{2*} \nabla_{\mathbf{t}^{1}} r_{i}^{2} \left(\mathbf{t}^{2*} \right).$

Using Lemma 4.3 and identifying corresponding multipliers we obtain

$$= \nabla_{\mathbf{t}^{1}} \mathfrak{f}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*})) - \sum_{i=1}^{ncstr_{1}} \mathfrak{w}_{i}^{1*} \nabla_{\mathbf{t}^{1}} r_{i}^{1}(\mathbf{t}^{1*}) - \sum_{i=1}^{ncstr_{2}} \mathfrak{w}_{i}^{2*} \nabla_{\mathbf{t}^{1}} r_{i}^{2}(\mathbf{t}^{2*})$$
$$= \nabla_{\mathbf{t}^{1}} L_{I}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}), \mathfrak{w}^{1*}, \mathfrak{w}^{2*}).$$

Analogously, $\mathbf{0} = \nabla_{\mathbf{t}^1} L_I(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}), \mathbf{w}^{1*}, \mathbf{w}^{2*})$. Together with the feasibility of the constraints and the complementarity (after identification of corresponding multipliers) this yields the first order optimality conditions of the full problem. The constraint qualifications carry over from the full problem since the constraints are unaltered.

Note that $\nabla_{\mathbf{A}} L_{I}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}), \mathbf{w}^{1*}, \mathbf{w}^{2*}) = \mathbf{0}$ due to the definition of $\mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*})$.

Hence $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt} (\mathbf{t}^{1*}, \mathbf{t}^{2*}), \mathbf{w}^{1*}, \mathbf{w}^{2*})$ is a critical point of the full problem. From the definition of the reduced problem we have

$$\mathfrak{f}\left(\mathbf{t}^{1*},\mathbf{t}^{2*},\mathbf{A}_{opt}\left(\mathbf{t}^{1*},\mathbf{t}^{2*}\right)\right)=f\left(\mathbf{t}^{1*},\mathbf{t}^{2*}\right)$$

The rest of the proof follows the ideas of Golub/Pereyra. For the sake of completeness we state the remaining steps.

Let $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$ be a global minimizer of the reduced problem on $\Omega_1 \times \Omega_2$ and let $\mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}) = \mathbf{B}_1(\mathbf{t}^{1*})^+ \mathbf{Z} (\mathbf{B}_2(\mathbf{t}^{2*})^+)^T$. Then $\mathfrak{f}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*})) = f(\mathbf{t}^{1*}, \mathbf{t}^{2*})$. Assume there exists $(\mathbf{t}^{1\dagger}, \mathbf{t}^{2\dagger}, \mathbf{A}^{\dagger})$ with $\mathbf{t}^{1\dagger} \in \Omega_1$, $\mathbf{t}^{2\dagger} \in \Omega_2$, such that $\mathfrak{f}(\mathbf{t}^{1\dagger}, \mathbf{t}^{2\dagger}, \mathbf{A}^{\dagger}) < \mathfrak{f}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}))$. For all $(\mathbf{t}^1, \mathbf{t}^2)$ we have $f(\mathbf{t}^1, \mathbf{t}^2) \leq \mathfrak{f}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A})$, hence

$$f\left(\mathbf{t}^{1\dagger},\mathbf{t}^{2\dagger}\right) \leq \mathfrak{f}\left(\mathbf{t}^{1\dagger},\mathbf{t}^{2\dagger},\mathbf{A}^{\dagger}\right) < \mathfrak{f}\left(\mathbf{t}^{1*},\mathbf{t}^{2*},\mathbf{A}_{opt}\left(\mathbf{t}^{1*},\mathbf{t}^{2*}\right)\right) = f\left(\mathbf{t}^{1*},\mathbf{t}^{2*}\right)$$

in contrast to the assumption. Thus $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}))$ is a global minimizer of the full problem in $\Omega_1 \times \Omega_2$.

(ii) Let $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$ be a global minimizer of the full problem for $(\mathbf{t}^1, \mathbf{t}^2) \in \Omega_1 \times \Omega_2$ and let $\mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*}) = \mathbf{B}_1(\mathbf{t}^{1*})^+ \mathbf{Z} (\mathbf{B}_2(\mathbf{t}^{2*})^+)^T$. Then there holds $f(\mathbf{t}^{1*}, \mathbf{t}^{2*}) \leq \mathfrak{f}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$. From the definition of the reduced functional one has $f(\mathbf{t}^{1*}, \mathbf{t}^{2*}) = \mathfrak{f}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*})) \leq \mathfrak{f}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$. Since $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$ is a global minimizer, equality holds, i.e.,

$$f\left(\mathbf{t}^{1*}, \mathbf{t}^{2*}\right) = \mathfrak{f}\left(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^{*}\right).$$

If there is an unique \mathbf{A}^* among all minimizing triplets of $\mathfrak{f}(\mathbf{t}^1, \mathbf{t}^2, \mathbf{A})$, it holds $\mathbf{A}^* = \mathbf{A}_{opt}(\mathbf{t}^{1*}, \mathbf{t}^{2*})$.

Let us assume that $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$ is *no* global minimizer of the reduced problem on $\Omega_1 \times \Omega_2$, i. e., there exist $(\mathbf{t}^{1\dagger}, \mathbf{t}^{2\dagger}) \in \Omega_1 \times \Omega_2$ with $f(\mathbf{t}^{1\dagger}, \mathbf{t}^{2\dagger}) < f(\mathbf{t}^{1*}, \mathbf{t}^{2*})$. Then, with $\mathbf{A}^{\dagger} = \mathbf{B}_1 (\mathbf{t}^{1\dagger})^+ \mathbf{Z} (\mathbf{B}_2 (\mathbf{t}^{2\dagger})^+)^T$, we have

$$f\left(\mathbf{t}^{1\dagger}, \mathbf{t}^{2\dagger}\right) = \mathfrak{f}\left(\mathbf{t}^{1\dagger}, \mathbf{t}^{2\dagger}, \mathbf{A}^{\dagger}\right) < f\left(\mathbf{t}^{1*}, \mathbf{t}^{2*}\right) = \mathfrak{f}\left(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^{*}\right)$$

in contrast to the assumption that $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$ is a global minimizer of the full problem.

From the argumentation one sees immediately that the statements carry over to the case of nonlinear equality constraints $\mathbf{s}^{1}(\mathbf{t}^{1}) = \mathbf{0}$ and $\mathbf{s}^{2}(\mathbf{t}^{2}) = \mathbf{0}$.

5 Bivariate tensor product splines with free knots.

After these preparations we can now apply the reduction technique to the full smoothing problem (3.10), (3.8). By continuity arguments analogously to the univariate case, see [27], [25, Theorem 4.2], we obtain

THEOREM 5.1 (EXISTENCE OF SOLUTIONS TO THE REDUCED SMOOTHING PROBLEM).

Let the set of feasible knots $\{(\mathbf{t}^1, \mathbf{t}^2) \in \mathbb{R}^{l_1} \times \mathbb{R}^{l_2} : \mathbf{C}_1 \mathbf{t}^1 - \mathbf{h}^1 \ge \mathbf{0}, \mathbf{C}_2 \mathbf{t}^2 - \mathbf{h}^2 \ge \mathbf{0}\}$ be nonempty, and let the following conditions be fulfilled for fixed $r_1 \in \{0, \ldots, q_1\}$, $0 \le q_1 < k_1$ and $r_2 \in \{0, \ldots, q_2\}$, $0 \le q_2 < k_2$:

- (C1) The knots satisfy $\tau_{j_1}^1 < \tau_{j_1+k_1-q_1}^1$ $(j_1 = q_1 + 1, \dots, n_1)$ and $\tau_{j_2}^2 < \tau_{j_2+k_2-q_2}^2$ $(j_2 = q_2 + 1, \dots, n_2).$
- (C2) The regularity conditions $m_1 \ge r_1$, $\mu_1 > 0$ and $m_2 \ge r_2$, $\mu_2 > 0$ are met.
- Then the reduced smoothing problem (3.11), (3.8) has a solution $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$.

We further have the smoothness of the reduced functional and, by applying Theorem 4.4, the equivalence of the full and reduced smoothing problem in the following sense:

THEOREM 5.2 (CORRESPONDENCE BETWEEN THE FULL AND REDUCED SMOOTH-ING PROBLEM).

Let $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$ be a feasible knot sequence, i. e.,

$$(\mathbf{t}^{1*}, \mathbf{t}^{2*}) \in \left\{ (\mathbf{t}^1, \mathbf{t}^2) \in \mathbb{R}^{l_1} imes \mathbb{R}^{l_2} \, : \, \mathbf{C}_1 \mathbf{t}^1 - \mathbf{h}^1 \ge \mathbf{0}, \mathbf{C}_2 \mathbf{t}^2 - \mathbf{h}^2 \ge \mathbf{0}
ight\},$$

and let the following conditions be fulfilled for fixed $r_1 \in \{0, ..., q_1\}, 0 \le q_1 < k_1$ and $r_2 \in \{0, ..., q_2\}, 0 \le q_2 < k_2$:

- (C1) The knots satisfy $\tau_{j_1}^1 < \tau_{j_1+k_1-q_1}^1$ $(j_1 = q_1 + 1, \dots, n_1)$ and $\tau_{j_2}^2 < \tau_{j_2+k_2-q_2}^2$ $(j_2 = q_2 + 1, \dots, n_2).$
- (C2) The regularity conditions $m_1 \ge r_1$, $\mu_1 > 0$ and $m_2 \ge r_2$, $\mu_2 > 0$ are met.
- (C3) The free knots $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$ are simple, and there holds $k_1 \geq 3$ and $k_2 \geq 3$.

Then the following relations hold for the full smoothing problem (3.10), (3.8) and the reduced smoothing problem (3.11), (3.8): The reduced function \mathbf{F} is smooth on the feasible set $\{(\mathbf{t}^1, \mathbf{t}^2) \in \mathbb{R}^{l_1} \times \mathbb{R}^{l_2} : \mathbf{C}_1 \mathbf{t}^1 - \mathbf{h}^1 \geq \mathbf{0}, \mathbf{C}_2 \mathbf{t}^2 - \mathbf{h}^2 \geq \mathbf{0}\}$.

- (a) If $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$ is a critical point (or a global minimizer) of (3.11), (3.8) and \mathbf{A}^* satisfies (3.6b) at the point $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$, then $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$ is a critical point (or a global minimizer) of (3.10), (3.8) and it holds $f(\mathbf{t}^{1*}, \mathbf{t}^{2*}) = f(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$.
- (b) If $(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$ is a global minimizer of (3.10), (3.8), then $(\mathbf{t}^{1*}, \mathbf{t}^{2*})$ is a global minimizer of (3.11), (3.8). It holds $f(\mathbf{t}^{1*}, \mathbf{t}^{2*}) = \mathfrak{f}(\mathbf{t}^{1*}, \mathbf{t}^{2*}, \mathbf{A}^*)$ and (3.6b).

The last two Theorems are straightforward generalizations of the univariate results. Note, however, that the expressions from Section 4 derived for the gradient and Jacobian are crucial for efficient numerical methods.

6 Numerical solution of the reduced problem.

After having shown the equivalence of the full and reduced problem in the sense of Theorem 5.2 we will now consider the numerical solution of the reduced problem.

The reduced problem is a nonlinear least squares problem in the variables \mathbf{t}^1 and \mathbf{t}^2 with linear inequality constraints. Formally the full problem resembles a separable least squares problem with multiple right hand sides. Such a problem can be expressed as $\frac{1}{2} \|\mathbf{Z} - \mathbf{B}_1(\mathbf{t}^1)\mathbf{A}\|_F^2 \to \min_{\mathbf{t}^1,\mathbf{A}}$, see [13] and [17].

can be expressed as $\frac{1}{2} \|\mathbf{Z} - \mathbf{B}_1(\mathbf{t}^1)\mathbf{A}\|_F^2 \to \min_{\mathbf{t}^1,\mathbf{A}}$, see [13] and [17]. A naive realization transforms the problem $\frac{1}{2} \|\mathbf{Z} - \mathbf{B}_1(\mathbf{t}^1)\mathbf{A}\mathbf{B}_2^T(\mathbf{t}^2)\|_F^2 \to \min$ into standard matrix-vector form $\frac{1}{2} \|\operatorname{vec}(\mathbf{Z}) - (\mathbf{B}_2(\mathbf{t}^2) \otimes \mathbf{B}_1(\mathbf{t}^1)) \operatorname{vec}(\mathbf{A})\|_2^2 \to \min$ and then applies the reduction technique. In this case we have to factorize the large matrix $\mathbf{B}_2 \otimes \mathbf{B}_1 \in \mathbb{R}^{(m_1 m_2) \times (n_1 n_2)}$ to compute the Jacobian of the reduced functional. However, a closer investigation of the structure shows that in every block of the Jacobian the same Fréchet derivative occurs. This structure has to be exploited for the solution process, in particular when solving large scale problems. In the case of separable least squares problems with multiple right hand sides this has first been done in [13].

The method of choice for solving the reduced problem is a generalized Gauss– Newton method. In every step one has to solve the quadratic model problem

$$\begin{split} \psi(\mathbf{t}^1 + \Delta \mathbf{t}^1, \mathbf{t}^2 + \Delta \mathbf{t}^2) &:= \\ \frac{1}{2} \left\| \mathbf{F}(\mathbf{t}^1, \mathbf{t}^2) + \partial_1 \mathbf{F}(\mathbf{t}^1, \mathbf{t}^2) [\Delta \mathbf{t}^1] + \partial_2 \mathbf{F}(\mathbf{t}^1, \mathbf{t}^2) [\Delta \mathbf{t}^2] \right\|_F^2 \to \min_{\Delta \mathbf{t}^1 \in \mathbb{R}^{l_1}, \Delta \mathbf{t}^2 \in \mathbb{R}^{l_2}} \end{split}$$

subject to the constraints

$$\mathbf{C}_1\mathbf{t}^1 + \mathbf{C}_1\Delta\mathbf{t}^1 - \mathbf{h}^1 \ge \mathbf{0}, \quad \mathbf{C}_2\mathbf{t}^2 + \mathbf{C}_2\Delta\mathbf{t}^2 - \mathbf{h}^2 \ge \mathbf{0}.$$

This yields

$$\begin{split} \psi &= \frac{1}{2} \left\| \mathbf{F} + \boldsymbol{\partial}_{1} \mathbf{F} \Delta \mathbf{t}^{1} + \boldsymbol{\partial}_{2} \mathbf{F} \Delta \mathbf{t}^{2} \right\|_{F}^{2} \\ &= \frac{1}{2} \left\| \mathbf{F} + \sum_{\kappa=1}^{l_{1}} \boldsymbol{\partial}_{1} \mathbf{F} [\mathbf{e}^{\kappa}] \Delta \mathbf{t}_{\kappa}^{1} + \sum_{\kappa=1}^{l_{2}} \boldsymbol{\partial}_{2} \mathbf{F} [\mathbf{e}^{\kappa}] \Delta \mathbf{t}_{\kappa}^{2} \right\|_{F}^{2} \\ &= \frac{1}{2} \left\| \operatorname{vec} \left(\mathbf{F} \right) + \sum_{\kappa=1}^{l_{1}} \operatorname{vec} \left(\boldsymbol{\partial}_{1} \mathbf{F} [\mathbf{e}^{\kappa}] \right) \Delta \mathbf{t}_{\kappa}^{1} + \sum_{\kappa=1}^{l_{2}} \operatorname{vec} \left(\boldsymbol{\partial}_{2} \mathbf{F} [\mathbf{e}^{\kappa}] \right) \Delta \mathbf{t}_{\kappa}^{2} \right\|_{2}^{2} \\ &= \frac{1}{2} \left\| \operatorname{vec} \left(\mathbf{F} \right) + \mathbf{J} \left(\begin{array}{c} \Delta \mathbf{t}^{1} \\ \Delta \mathbf{t}^{2} \end{array} \right) \right\|_{2}^{2} \end{split}$$

with

$$\mathbf{J} := \begin{bmatrix} | & | & | & | \\ \operatorname{vec} \left(\boldsymbol{\partial}_1 \mathbf{F}[\mathbf{e}^1] \right) \cdots \operatorname{vec} \left(\boldsymbol{\partial}_1 \mathbf{F}[\mathbf{e}^{l_1}] \right) & \operatorname{vec} \left(\boldsymbol{\partial}_2 \mathbf{F}[\mathbf{e}^1] \right) \cdots \operatorname{vec} \left(\boldsymbol{\partial}_2 \mathbf{F}[\mathbf{e}^{l_2}] \right) \\ | & | & | & | & | \end{bmatrix}$$

The Jacobian $\mathbf{J} \in \mathbb{R}^{(m_1+n_1-r_1)(m_2+n_2-r_2)\times(l_1+l_2)}$ in the smoothing case (resp. $\mathbf{J} \in \mathbb{R}^{(m_1m_2)\times(l_1+l_2)}$ in the approximation case), which is in general full, can be computed column-wise.

When computing the Jacobian in the above way, the required QR factorizations of \mathbf{B}_1 and \mathbf{B}_2 for $\partial_1 \mathbf{F}$ and $\partial_2 \mathbf{F}$ have to be computed only once and can then by applied to different right hand sides. All algorithms for the computation of the Jacobian—including the use of the Kaufman approximation and the exploitation of sparsity—carry over from the univariate case. For example, in the case of spline approximation we obtain the Jacobian

$$\boldsymbol{\partial}_{1}\mathbf{F}(\mathbf{t}^{1},\mathbf{t}^{2})[\Delta\mathbf{t}^{1}] = -\left\{\mathbf{P}_{B_{1}}^{\perp}(\boldsymbol{\partial}_{1}\mathbf{B}_{1}[\Delta\mathbf{t}^{1}])\mathbf{B}_{1}^{+} + \left(\mathbf{P}_{B_{1}}^{\perp}(\boldsymbol{\partial}_{1}\mathbf{B}_{1}[\Delta\mathbf{t}^{1}])\mathbf{B}_{1}^{+}\right)^{T}\right\}\mathbf{Z}\mathbf{P}_{B_{2}}$$

and the Kaufman approximation

$$\mathbf{J}_{K}(\mathbf{t}^{1},\mathbf{t}^{2})[\Delta\mathbf{t}^{1}] = -\mathbf{P}_{B_{1}}^{\perp}(\boldsymbol{\partial}_{1}\mathbf{B}_{1}[\Delta\mathbf{t}^{1}])\mathbf{B}_{1}^{+}\mathbf{Z}\mathbf{P}_{B_{2}}.$$

This procedure bears great resemblance to the computation of the Jacobian for separable least squares problems with multiple right hand sides. If we set $\mathbf{B}_2(\mathbf{t}^2) = \mathbf{I}$ we arrive straightforward at the results of Golub/LeVeque as a special case. The exploitation of structure for these problems has been intensively investigated during the last years, see [17], [18], [11], [28]. Kaufman/Sylvester report on a drastic reduction in computing time for real world problems with thousands of parameters and millions of observations. While in this case the linear least squares problems were full, Soo/Bates [28] investigate the additional sparsity of the observation matrix. As one example they consider self-modeling free-knot splines, but only in the univariate and not in the tensor product case.

In conclusion, it can be stated that the algorithms for the computation of the Jacobian and the generalized Gauss–Newton method developed in [27], [25] for the univariate case carry over to bivariate free knot splines, in principle.

7 Numerical tests.

For the computation of bivariate tensor product splines with free knots we have implemented a method for the solution of the reduced problem in MATLAB. We used the code FMINCON from the MATLAB Optimization Toolbox [5] and the code NPSOL [12] on a PENTIUM 1 GHz processor. Both methods are SQPmethods which use a BFGS-update of the Hessian of the Lagrangian. Gradients have been computed via finite differences, but exact gradients are also available using the formulae given in [25]. In all examples we took $a_1 = x_1$, $b_1 = x_{m_1}$ and $a_2 = y_1$, $b_2 = y_{m_2}$.

7.1 Bivariate Titanium Heat Data.

In our first example we consider the well-known Titanium Heat Data [2]. Building the tensor product of the univariate data, i.e., $z_{i_1,i_2} = y_{i_1} \times y_{i_2}$, we

get $(m_1 = 49) \times (m_2 = 49)$ points in $[595, 1075] \times [595, 1075]$, see Figure 7.1. We want to approximate these 2401 data points by $n_1 = 11$ cubic B-splines in *x*-direction and $n_2 = 10$ cubic B-splines in *y*-direction. Considering all inner knots as free we achieve at $l_1 = 7$ and $l_2 = 6$.



Figure 7.1: Bivariate Titanium Heat Data: data points



Figure 7.2: Bivariate Titanium Heat Data: spline s, initial knot sequence

If we choose equidistant interior knots we obtain the approximation shown in Figure 7.2. Besides the oscillations in the flat part one observes that, in particular, the approximation near the peak is very poor. Optimizing the location of the knots the oscillations disappear and the residual decreases to 5%. The results are summarized in Table 7.1. Note that the MATLAB procedure cannot achieve the desired accuracy. The resulting spline after the optimization with NPSOL is shown in Figure 7.3. In Figure 7.4 we plot the location of the knots and the corresponding contour before and after the optimization.

	initial knot sequence	FMINCON	NPSOL
$\ \mathbf{F}\ $	$9.777865 \mathrm{E}{+}00$	$6.652674 ext{ E-01}$	$5.057175 ext{ E-01}$
steps		87	56
function calls		1307	550
time [sec]		8.22	8.63
return code		max. no. iterations	successful

Table 7.1: Bivariate Titanium Heat Data: comparison of FMINCON and NPSOL



Figure 7.3: Bivariate Titanium Heat Data: optimized location of knots, NPSOL

Above example clarifies the importance of a good knot sequence. Although the residuals of both approximations do not differ by several magnitudes the approximating spline with equidistant knots is, in fact, unusable due to its high oscillations. By minimizing the least square error with respect to the free knots not only the approximation error gets smaller, even the approximating spline



Figure 7.4: Bivariate Titanium Heat Data: contour plot and knots before and after the optimization

itself becomes visually more pleasant.

7.2 EOS Aluminium Data.

In a second example we use a standard data set for bivariate *constrained* approximation, see e.g. [4]. The $(m_1 = 10) \times (m_2 = 6)$ data points in $[-0.07, 1.13] \times [-2.3, 0]$ describe an equation of state (EOS) for aluminium. Represented is pressure as a function of density and temperature on a log-log scale. The data are in monotone position, see Figure 7.5.

Although this data set is relatively small it is difficult to approximate. We use $n_1 = 8$ quadratic B-splines in x-direction, $n_2 = 5$ quadratic B-splines in y-direction and the smoothing parameters $\mu_1 = \mu_2 = 1.0 \text{ E-}08$, $r_1 = r_2 = 2$. For equidistant inner knots we have the unsatisfying approximation shown in Figure 7.6.

Optimizing the location of knots one obtains, for example with FMINCON, the spline shown in Figure 7.7. The essential oscillations have disappeared. Table 7.2 and Figure 7.8 summarize the results. The resulting approximating

	initial knot sequence	FMINCON	NPSOL
$\ \mathbf{F}\ $	$1.587922 \mathrm{E}{+}01$	$1.228879 \to +00$	$1.027007\mathrm{E}{+}00$
steps		65	36
function calls		603	691
time [sec]		2.03	2.71
return code		successful	succesful

Table 7.2: EOS Aluminium Data: comparison of FMINCON and NPSOL



Figure 7.5: EOS Aluminium Data: data set



Figure 7.6: EOS Aluminium Data: spline $\boldsymbol{s},$ initial knot sequence



Figure 7.7: EOS Aluminium Data: optimized knots, FMINCON

spline is almost monotone, it holds min $s_y \approx -12$, max $s_y \approx 86!$ Note that in this example both methods abort in the case of pure spline approximation without smoothing ($\mu_1 = \mu_2 = 0$) since rank deficient observation matrices—hence loss of differentiability—do occur in the optimization process.



Figure 7.8: EOS Aluminium Data: contour plot and knots before and after the optimization

The almost-monotonicity suggests a further area of application for bivariate free knot splines: In so-called fit-and-modify methods for constrained interpola-

T. SCHÜTZE AND H. SCHWETLICK

tion one has to provide good estimates for derivatives. The parameters of the constrained spline are then computed in such a way that the resulting spline deviates as small as possible from the given spline, but satisfies the shape constraints. Since the derivatives of a bivariate free knot spline are of good quality, in general, they qualify as starting parameters for fit-and-modify methods. Figure 7.9 represents a monotone interpolating spline that has been computed using a special fit-and-modify interpolation method proposed in [23] by Schmidt/Bastian-Walther.

approximation type	least squares approx.	constrained interpolation
type of optimization problem	nonlinear	quadratic
constraints on derivatives	unconstrained	monotonicity
time [sec]	2.03	45.62
$\min s_y$	-12	0
$\max s_u$	86	905

86

Table 7.3: Comparison of our method with the fit-and-modify interpolation method [23] by Schmidt/Walther



Figure 7.9: EOS Aluminium Data: monotone spline, fit-and-modify interpolation method [23] by Schmidt/Walther

176

 $\max s_y$

Table 7.3 compares both methods using the EOS Aluminium Data. By combining both methods we expect substantial advantages since the unconstrained free knot least squares spline will provide good initial derivative estimates for the subsequent fit-and-modify iteration method so an essential reduction of computing time can be expected.

REFERENCES

- Å. Björck, Numerical Methods for Least Squares Problems, SIAM, Philadelphia, PA, 1996.
- 2. C. de Boor, A Practical Guide to Splines, Springer-Verlag, New York, revised ed., 2001.
- K. Brenner, Bivariate Quadratmittelapproximation mittels glättender Tensorprodukt-B-Splines (Bivariate least squares approximation by smoothing tensor product B-splines), Diplomarbeit, Martin-Luther-Universität, Halle-Wittenberg, 1990.
- R. E. Carlson and F. N. Fritsch, Monotone piecewise bicubic interpolation, SIAM J. Numer. Anal., 22 (1985), pp. 386–400.
- 5. T. Coleman, M. A. Branch, and A. Grace, *Optimization Toolbox User's Guide*, *Version 2*, The MathWorks, Inc., 1999.
- P. Dierckx, An algorithm for surface-fitting with spline functions, IMA J. Numer. Anal., 1 (1981), pp. 267–283.
- P. Dierckx, A fast algorithm for smoothing data on a rectangular grid while using spline functions, SIAM J. Numer. Anal., 19 (1982), pp. 1286–1305.
- P. Dierckx, *FITPACK user guide, part 2: Surface fitting routines*, Tech. Rep. TW Report 122, Department of Computer Science, Katholieke Universiteit, Leuven, Belgium, 1989.
- S. Ewald, H. Mühlig, and B. Mulansky, *Bivariate interpolating and smoothing tensor product splines*, in Splines in Numerical Analysis, Proceedings ISAM-89, J. W. Schmidt and H. Späth, eds., Vol. 52 of Mathematical Research, Akademie-Verlag, Berlin, 1989, pp. 55–68.
- D. W. Fausett and C. T. Fulton, Large least squares problems involving Kronecker products, SIAM J. Matrix Anal. Appl., 15 (1994), pp. 219–227.
- D. M. Gay and L. Kaufman, Tradeoffs in algorithms for separable nonlinear least squares, in Computational and Applied Mathematics, I. Algorithms and Theory, Selected Papers IMACS 13th World Congress, Dublin, Ireland, 1991, 1992, pp. 179– 183.
- P. E. Gill, W. Murray, M. A. Saunders, and M. H. Wright, User's guide for NPSOL (version 4.0): A Fortran package for nonlinear programming, Tech. Report SOL 86-2, Department of Operations Research, Standford University, 1986.
- G. H. Golub and R. J. LeVeque, Extensions and uses of the variable projection algorithm for solving nonlinear least squares problems, in Proceedings 1979 Army Numerical Analysis and Computer Science Conference, Army Research Office Report 79–3, 1979, pp. 1–12.
- G. H. Golub and V. Pereyra, The differentiation of pseudoinverses and nonlinear least squares problems whose variables separate, SIAM J. Numer. Anal., 10 (1973), pp. 413–432.

- C. L. Hu and L. L. Schumaker, *Bivariate natural spline smoothing*, in Delay Equations, Approximation and Applications, Mannheim 1994, G. Meinardus and G. Nürnberger, eds., Vol. 74 of ISNM, Birkhäuser, 1985, pp. 165–179.
- C. L. Hu and L. L. Schumaker, *Complete spline smoothing*, Numer. Math., 49 (1986), pp. 1–10.
- L. Kaufman and G. S. Sylvester, Separable nonlinear least squares with multiple right-hand sides, SIAM J. Matrix Anal. Appl., 13 (1992), pp. 68–89.
- L. Kaufman, G. S. Sylvester, and M. H. Wright, Structured linear least-squares problems in system identification and separable nonlinear data fitting, SIAM J. Optim., 4 (1994), pp. 847–871.
- G. Meinardus, G. Nürnberger, and G. Walz, Bivariate segment approximation and splines, Adv. Comput. Math., 6 (1996), pp. 25–45.
- B. Mulansky, Glättung mittels zweidimensionaler Tensorprodukt-Splinefunktionen, Wiss. Z. Tech. Univ. Dresden, 39 (1990), pp. 187–190.
- 21. G. Nürnberger, Bivariate segment approximation and free knot splines: Research problems 96-4, Constr. Approx., 12 (1996), pp. 555–558.
- 22. T. Pigorsch, Bivariate Quadratmittelapproximation unter Verwendung von Tensorprodukt-B-Splines im Falle von Rechteckgitterdaten (Bivariate least squares approximation for gridded data using tensor product B-splines), Diplomarbeit, Martin-Luther-Universität Halle-Wittenberg, 1991.
- J. W. Schmidt and M. Walther, Gridded data interpolation with restrictions on the first order derivatives, in Multivariate Approximation and Splines, G. Nürnberger, J. W. Schmidt, and G. Walz, eds., Vol. 125 of ISNM, Birkhäuser, Basel, 1997, pp. 291–307.
- 24. T. Schütze, Diskrete Quadratmittelapproximation durch Splines mit freien Knoten (Discrete least squares approximation by splines with free knots), Dissertation, Technische Universität Dresden, 1997.
- T. Schütze and H. Schwetlick, Constrained approximation by splines with free knots, BIT, 37 (1997), pp. 105–137.
- H. Schwetlick and V. Kunert, Spline smoothing under constraints on derivatives, BIT, 33 (1993), pp. 512–528.
- 27. H. Schwetlick and T. Schütze, Least squares approximation by splines with free knots, BIT, 35 (1995), pp. 361–384.
- Y. W. Soo and D. M. Bates, *Loosely coupled nonlinear least squares*, Computational Statistics and Data Analysis, 14 (1992), pp. 249–259.